

- 2 Beets GL, Figueiredo NF, Beets-Tan RG. Management of rectal cancer without radical resection. *Annu Rev Med* 2017; **68**: 169–82.
- 3 Rullier E, Rouanet P, Tuech J-J, et al. Organ preservation for rectal cancer (GRECCAR 2): a prospective, randomised, open-label, multicentre, phase 3 trial. *Lancet* 2017; published online June 7. [http://dx.doi.org/10.1016/S0140-6736\(17\)31056-5](http://dx.doi.org/10.1016/S0140-6736(17)31056-5).
- 4 Verseveld M, de Graaf EJ, Verhoef C, et al. Chemoradiation therapy for rectal cancer in the distal rectum followed by organ-sparing transanal endoscopic microsurgery (CARTS study). *Br J Surg* 2015; **102**: 853–60.
- 5 Pucciarelli S, De Paoli A, Guerrieri M, et al. Local excision after preoperative chemoradiotherapy for rectal cancer: results of a multicenter phase II clinical trial. *Dis Colon Rectum* 2013; **56**: 1349–56.
- 6 Relton C, Torgerson D, O’Cathain A, Nicholl J. Rethinking pragmatic randomised controlled trials: introducing the “cohort multiple randomised controlled trial” design. *BMJ* 2010; **340**: c1066.
- 7 Beets GL, Figueiredo NL, Habr-Gama A, van de Velde CJ. A new paradigm for rectal cancer: organ preservation: introducing the International Watch & Wait Database (IWW). *Eur J Surg Oncol* 2015; **41**: 1562–64.

## Post-partum depression—a glimpse of light in the darkness?

Published Online

June 12, 2017

[http://dx.doi.org/10.1016/S0140-6736\(17\)31546-5](http://dx.doi.org/10.1016/S0140-6736(17)31546-5)

See [Articles](#) page 480

There can be little doubt about the importance of mood episodes in pregnancy and following childbirth.<sup>1</sup> Mood episodes are common—post-partum depression is the most common medical complication of maternity, affecting around one in ten new mothers.<sup>2</sup> They can also be severe—episodes of post-partum psychosis represent some of the most serious episodes of illness seen in psychiatry.<sup>3</sup> Perinatal mood episodes cause substantial impairment to women and have a wide ranging impact on their babies, families, and society. The total long-term cost to society of perinatal depression, anxiety, and psychosis has been estimated to be £8.1 billion for each 1-year cohort of births in the UK.<sup>4</sup>

Despite the clear need, there are limited options for the management of these disorders. In addition to psychological approaches such as cognitive behavioural therapy, medications, particularly for more severe episodes of illness, are a mainstay of treatment.<sup>5</sup> Because, at least in part, pregnant and breastfeeding women are excluded from clinical trials, decisions about

medication for perinatal mood episodes are difficult; they must be made by extrapolating the data available for their use at other times of women’s lives. It is clear, therefore, that developing new, evidence-based treatments is essential.

In *The Lancet*, Stephen Kaner and colleagues<sup>6</sup> report the results of a double-blind, randomised, placebo-controlled, phase 2 study of a new treatment for post-partum depression. The compound, brexanolone, is an intravenous formulation of allopregnanolone, a positive allosteric modulator of  $\gamma$ -aminobutyric acid (GABA) receptors. In this small study in 21 women with severe post-partum depression, infusion of brexanolone resulted in a rapid, sustained, statistically significant, and clinically meaningful response compared with placebo (at 60 h, mean reduction in Hamilton Rating Scale for Depression [HAM-D] total score from baseline 21.0 points [SE 2.9] in the brexanolone group vs 8.8 points [SE 2.8] in the placebo group; difference -12.2, 95% CI -3.67 to -20.77;  $p=0.0075$ ; effect size 1.2). There were significant differences between groups from 24 h; by 60 h, seven (70%) women in the brexanolone group had achieved remission compared with one (9%) woman in the placebo group ( $p=0.0364$ ). These results, while based on a small sample, are very impressive; indeed, some readers might feel that they are too good to be true. For women with post-partum depression and the professionals who treat them, however, these findings are promising.

The investigators are to be applauded for targeting women with severe episodes of post-partum depression, and for showing that a clinical trial in women with this condition is feasible.

The current study, it must be noted, involved only ten women treated with brexanolone. The pressing



Science Photo Library C031/2967

need is therefore for replication of these results in larger phase 3 trials. Further questions may then need to be addressed that are raised but not answered by this study. It does not tell us, for example, if this is a treatment for post-partum episodes specifically, or for depression more generally? Is the treatment effective in other psychiatric conditions triggered by childbirth such as post-partum psychosis or in a wider group of reproductive and endocrine-related mood disorders such as those related to menstruation and menopause?

The findings have potentially important implications for our understanding of the pathophysiology of post-partum mood disorders and, given what is known about the action of brexanolone, provide further evidence implicating neuroactive steroids in general, and the GABA type A (GABA<sub>A</sub>) receptor  $\delta$  subunit in particular. Other research disciplines, such as neuroimaging and genetics, can further explore this promising avenue of research.

The need for a 60 h intravenous infusion with brexanolone, although possibly not an issue for women with severe post-partum depression, could be problematic if the treatment is found to be effective in less severe forms of the disorder. To this end, it will be interesting to see if it will be possible to develop GABA<sub>A</sub> positive allosteric modulators that can be administered orally with similar efficacy. Finally, in addition to the treatment of women who are currently

symptomatic, will this or similar treatments be suitable for the prevention of episodes in women at high risk?

Those of us hoping for the development of effective pharmacological treatments that specifically target post-partum depression have, like our patients, felt in a dark place. With the very encouraging results of this trial, perhaps we can begin to see the first glimpses of light.

Ian Jones

National Centre for Mental Health, MRC Centre for Neuropsychiatric Genetics and Genomics, Cardiff University, Cardiff, CF24 4HQ, UK  
Jonesir1@cf.ac.uk

I declare no competing interests.

- 1 Howard LM, Piot P, Stein A. No health without perinatal mental health. *Lancet* 2014; **384**: 1723–24.
- 2 Howard LM, Molyneaux E, Dennis CL, Rochat T, Stein A, Milgrom J. Non-psychotic mental disorders in the perinatal period. *Lancet* 2014; **384**: 1775–88.
- 3 Jones I, Chandra PS, Dazzan P, Howard LM. Bipolar disorder, affective psychosis, and schizophrenia in pregnancy and the post-partum period. *Lancet* 2014; **384**: 1789–99.
- 4 Bauer A, Parsonage M, Knapp M, Lemmi V, Adelaja B. Costs of perinatal mental health problems. London: London School of Economics and Political Science, 2014. <http://eprints.lse.ac.uk/59885/> (accessed May 4, 2017).
- 5 National Institute for Health and Care Excellence. Antenatal and postnatal mental health: clinical management and service guidance. Clinical guideline [CG192]. June, 2015. <https://www.nice.org.uk/guidance/cg192> (accessed May 4, 2017).
- 6 Kanes S, Colquhoun H, Gunduz-Bruce H, et al. Brexanolone (SAGE-547 injection) in post-partum depression: a randomised controlled trial. *Lancet* 2017; published online June 12. [http://dx.doi.org/10.1016/S0140-6736\(17\)31264-3](http://dx.doi.org/10.1016/S0140-6736(17)31264-3).

## Preventing major gastrointestinal bleeding in elderly patients



Antiplatelet therapy is the most frequently recommended treatment to prevent recurrent ischaemic events in patients who have had an ischaemic stroke, an acute coronary syndrome, or symptomatic peripheral arterial disease. The most frequently used drugs are aspirin or clopidogrel. Most guidelines recommend lifelong intake of antiplatelet therapy. However, randomised trials that have investigated the benefit of antiplatelet therapy had an observation period of between 2 years and 4 years.<sup>1</sup> Therefore, we lack data on the long-term benefit and risk of antiplatelet therapy across long time periods, particularly in elderly patients.

In *The Lancet*, Linxin Li and colleagues<sup>2</sup> report bleeding events and outcomes in 3166 patients with

first transient ischaemic attack, ischaemic stroke, or myocardial infarction who were treated with antiplatelet drugs (mainly aspirin) and were followed prospectively for 10 years. Half of the patients (n=1582) were aged 75 years or older.

Major bleeding and fatal bleeding were significantly related to age and showed a steep increase in incidence above the age of 75 years. The hazard ratio for major upper gastrointestinal bleeds was 4.13 for age 75 years or older and 10.26 for those bleeds that were disabling or fatal. The proportion of gastrointestinal bleeding events that were disabling or fatal was higher than the proportion of ischaemic stroke or intracerebral haemorrhage. At age 75 years or older, most major upper gastrointestinal

Published Online  
June 13, 2017  
[http://dx.doi.org/10.1016/S0140-6736\(17\)31507-6](http://dx.doi.org/10.1016/S0140-6736(17)31507-6)  
See [Articles](#) page 490